U.S. DEPARTMENT OF COMMERCE
National Technical Information Service

PB-284 671

# Oxbow Reservoir Baker County, Oregon, and Adams County, Idaho

National Eutrophication Survey, Corvallis, Oreg

Prepared for

Corvallis Environmental Research Lab, Oreg

Mar 78

PB 284 671

# U.S. ENVIRONMENTAL PROTECTION AGENCY
# NATIONAL EUTROPHICATION SURVEY

## WORKING PAPER SERIES

CORVALLIS ENVIRONMENTAL RESEARCH LABORATORY - CORVALLIS, OREGON

and

ENVIRONMENTAL MONITORING & SUPPORT LABORATORY - LAS VEGAS, NEVADA

REPRODUCED BY
NATIONAL TECHNICAL
INFORMATION SERVICE
U. S. DEPARTMENT OF COMMERCE
SPRINGFIELD, VA. 22161

G.P.O. 699-440

## TECHNICAL REPORT DATA
*(Please read Instructions on the reverse before completing)*

| 1. REPORT NO. | 2. | 3. RECIPIENT'S ACCESSION NO. PB284671 |
|---|---|---|
| 4. TITLE AND SUBTITLE<br>Report on Oxbow Reservoir, Baker County, Oregon, Adams County, Idaho | | 5. REPORT DATE<br>Prepared March 1978<br>6. PERFORMING ORGANIZATION CODE |
| 7. AUTHOR(S)<br>Anonymous (National Eutrophication Survey Staff) | | 8. PERFORMING ORGANIZATION REPORT NO.<br>Working Paper # 832 |
| 9. PERFORMING ORGANIZATION NAME AND ADDRESS<br>Pacific Northwest Environmental Research Laboratory<br>U.S. Environmental Protection Agency<br>200 S. W. 35th St.<br>Corvallis, Oregon 97330 | | 10. PROGRAM ELEMENT NO.<br>1BA029<br>11. CONTRACT/GRANT NO. |
| 12. SPONSORING AGENCY NAME AND ADDRESS<br>Office of Research and Development<br>U.S. Environmental Protection Agency<br>401 M Street, S. W.<br>Washington, DC 20460 | | 13. TYPE OF REPORT AND PERIOD COVERED<br>Final - 1975<br>14. SPONSORING AGENCY CODE |

15. SUPPLEMENTARY NOTES
Prepared in cooperation with the U.S. Environmental Protection Agency, National Environmental Research Center, Las Vegas, Nevada 89114

16. ABSTRACT
Annual total phosphorus and total nitrogen loadings to the lake were estimated and subdivided according to either point or non-point source origin. An assessment of the lake's trophic condition and limiting nutrient is also provided. All data collected by the U.S.E.P.A. National Eutrophication Survey during the one year study of the lake and its tributaries are included in the report.

17. KEY WORDS AND DOCUMENT ANALYSIS

| a. DESCRIPTORS | b. IDENTIFIERS/OPEN ENDED TERMS | c. COSATI Field/Group |
|---|---|---|
| eutrophication<br>lakes<br>nutrients<br>phosphorus<br>nitrogen<br>loading | National Eutrophication Survey<br>Lake Survey | 05B<br>05C |

| 18. DISTRIBUTION STATEMENT | 19. SECURITY CLASS *(This Report)* | 21. NO. OF PAGES |
|---|---|---|
| Release Unlimited | Unclassified | 39 |
| | 20. SECURITY CLASS *(This page)* | 22. PRICE |
| | | A03-A01 |

EPA Form 2220-1 (Rev. 4-77) PREVIOUS EDITION IS OBSOLETE

# REPORT
# ON
# OXBOW RESERVOIR
# BAKER COUNTY, OREGON
# ADAMS COUNTY, IDAHO
# EPA REGION X
# WORKING PAPER No. 832

WITH THE COOPERATION OF THE
OREGON DEPARTMENT OF ENVIRONMENTAL QUALITY,
THE IDAHO DEPARTMENT OF HEALTH AND WELFARE,
AND THE OREGON NATIONAL GUARD

MARCH, 1978

CONTENTS


| | Page |
|---|---|
| Foreword | ii |
| List of Oregon Lakes and Reservoirs | iv |
| Lake or Reservoir Drainage Area Map | v |
| **Sections** | |
| I. Conclusions | 1 |
| II. Drainage Basin Characteristics | 4 |
| III. Water Quality Summary | 5 |
| IV. Nutrient Loadings | 9 |
| V. Literature Reviewed | 13 |
| VI. Appendices | 14 |

# FOREWORD

The National Eutrophication Survey was initiated in 1972 in response to an Administration commitment to investigate the nationwide threat of accelerated eutrophication to freshwater lakes and reservoirs.

## OBJECTIVES

The Survey was designed to develop, in conjunction with state environmental agencies, information on nutrient sources, concentrations, and impact on selected freshwater lakes as a basis for formulating comprehensive and coordinated national, regional, and state management practices relating to point-source discharge reduction and non-point source pollution abatement in lake watersheds.

## ANALYTIC APPROACH

The mathematical and statistical procedures selected for the Survey's eutrophication analysis are based on related concepts that:

a. A generalized representation or model relating sources, concentrations, and impacts can be constructed.

b. By applying measurements of relevant parameters associated with lake degradation, the generalized model can be transformed into an operational representation of a lake, its drainage basin, and related nutrients.

c. With such a transformation, an assessment of the potential for eutrophication control can be made.

## LAKE ANALYSIS

In this report, the first stage of evaluation of lake and watershed data collected from the study lake and its drainage basin is documented. The report is formatted to provide state environmental agencies with specific information for basin planning [§303(e)], water quality criteria/standards review [§303(c)], clean lakes [§314(a,b)], and water quality monitoring [§106 and §305(b)] activities mandated by the Federal Water Pollution Control Act Amendments of 1972.

Beyond the single lake analysis, broader based correlations between nutrient concentrations (and loading) and trophic condition are being made to advance the rationale and data base for refinement of nutrient water quality criteria for the Nation's fresh water lakes. Likewise, multivariate evaluations for the relationships between land use, nutrient export, and trophic condition, by lake class or use, are being developed to assist in the formulation of planning guidelines and policies by EPA and to augment plans implementation by the states.

ACKNOWLEDGEMENT

The staff of the National Eutrophication Survey (Office of Research & Development, U. S. Environmental Protection Agency) expresses sincere appreciation to the Oregon Department of Environmental Quality and the Idaho Department of Health and Welfare for professional involvement and to the Oregon National Guard for conducting the tributary sampling phase of the Survey.

William H. Young, Director; Harold L. Sawyer, Administrator, and the staff of the Water Quality Control Division of the Oregon Department of Environmental Quality; and the staff of the Division of Environment of the Idaho Department of Health and Welfare provided invaluable lake documentation and counsel during the Survey, reviewed the preliminary reports, and provided critiques most useful in the preparation of this Working Paper series.

Brigadier General Richard A. Miller, the Adjutant General of Oregon, and Project Officer Lt. Colonel John Mewha, who directed the volunteer efforts of the Oregon National Guardsmen, are also gratefully acknowledged for their assistance to the Survey.

NATIONAL EUTROPHICATION SURVEY

STUDY LAKES and RESERVOIRS

STATE OF OREGON

| NAME | COUNTY |
|---|---|
| Brownlee | Baker, OR; Washington, ID |
| Diamond | Douglas |
| Hells Canyon | Baker, Wallowa, OR; Adams, Idaho, ID |
| Hills Creek | Lane |
| Owyhee | Malhuer |
| Oxbow | Baker, OR; Adams, ID |
| Suttle | Jefferson |
| Waldo | Lane |


116°55'
116°50'
SNAKE
RIVER
06A1/03A2
outlet
01 dam
OXBOW RESERVOIR
Tributary Sampling Site
Lake Sampling Site
0 1 2 3 4 5 Km.
0 1 2 3 Mi.
Scale
Salt
Creek
02
OREGON
IDAHO
44°55'
Ore.
Map Location
RESERVOIR
OXBOW
River
03
06B1
Wildhorse
06A2
Brownlee Dam
Brownlee
Reservoir
44°50'

OXBOW RESERVOIR

STORET NO. 4106

# I. CONCLUSIONS

## A. Trophic Condition:

Survey data indicate Oxbow Reservoir is eutrophic. It ranked sixteenth in overall trophic quality when the 21 Oregon and Idaho lakes and reservoirs sampled in 1975 were compared using a combination of six water quality parameters*. Seventeen of the water bodies had less median total phosphorus and dissolved orthophosphorus, all of the others had less median inorganic nitrogen, 13 had less mean chlorophyll a, and 14 had greater mean Secchi disc transparency. Some depression of dissolved oxygen with depth occurred at stations 1 and 2 in August and September.

Survey limnologists did not observe surface concentrations of algae; however, the relatively high chlorophyll a concentration at station 2 in April is indicative of a near-bloom condition (see page 7).

## B. Rate-Limiting Nutrient:

The algal assay results are not considered representative of conditions in the reservoir at the time the samples were collected (09/15/75).

The reservoir data indicate phosphorus limitation in April but nitrogen limitation in August and September.

---

* See Appendix A.

C. Nutrient Controllability:

1. Point sources--No known point sources directly impacted Oxbow Reservoir during the sampling year. However, those point sources contributing nutrients to upstream Brownlee Reservoir* probably affect Oxbow Reservoir as well, since 54% of the phosphorus input to Brownlee Reservoir was not retained and entered Oxbow Reservoir.

The sampling year phosphorus loading of 183.51 $g/m^2$ is over 20 times that proposed by Vollenweider (Vollenweider and Dillon, 1974) as a eutrophic loading (see page 12). Although Vollenweider's model may not be applicable to water bodies with short hydraulic retention times, the trophic condition of Oxbow Reservoir is evidence of excessive nutrient loads. Any improvement in water quality would require minimization of the nutrient loads from both non-point sources and the point sources impacting upstream Brownlee Reservoir.

2. Non-point sources--As considered in this report, "non-point" sources include those contributing nutrients as a result of human activities (e.g., agricultural practices, urbanization, extractive processes, etc.) as well as natural sources (i.e., direct precipitation, leaf fall, wild birds and animals, etc.).

The apparent non-point source phosphorus contribution accounted for all of the total phosphorus load during the sampling year. However, as stated above, part of this load is attributable to point

* Working Paper No. 827.

sources further upstream.

The Snake River contributed 99.0% of the total phosphorus load, and the Wildhorse River contributed 0.6%. The ungaged minor tributaries and immediate drainage contributed an estimated 0.4%.

## II. RESERVOIR AND DRAINAGE BASIN CHARACTERISTICS†

### A. Morphometry††:

1. Surface area: 5.66 kilometers².
2. Mean depth: 11.4 meters.
3. Maximum depth: 38.1 meters.
4. Volume: $64.758 \times 10^6$ m³.
5. Mean hydraulic retention time: 1 day.

### B. Tributary and Outlet:
(See Appendix C for flow data)

1. Tributaries -

| Name | Drainage area (km²)* | Mean flow (m³/sec)* |
|---|---|---|
| Snake River | 188,033.9 | 526.270 |
| Wildhorse River | 458.4 | 4.920 |
| Minor tributaries & immediate drainage - | 312.9 | 1.470 |
| Totals | 188,805.2 | 532.660** |

2. Outlet -

| | | |
|---|---|---|
| Snake River | 188,810.9*** | 532.660 |

### C. Precipitation****:

1. Year of sampling: 27.0 centimeters.
2. Mean annual: 27.3 centimeters.

---

† Table of metric equivalents--Appendix B.
†† McHugh, 1972; Martin and Hanson, 1966.
* For limits of accuracy, see Working Paper No. 175, "...Survey Methods, 1973-1976".
** Sum of inflows adjusted to equal outflow.
*** Includes area of reservoir.
**** See Working Paper No. 175.

## III. WATER QUALITY SUMMARY

Oxbow Reservoir was sampled three times during the open-water season of 1975 by means of a pontoon-equipped Huey helicopter. Each time, samples for physical and chemical parameters were collected from a number of depths at three stations on the reservoir (see map, page v). During each visit, a single depth-integrated (4.6 m to surface) sample was composited from the stations for phytoplankton identification and enumeration; and during the September visit, a single 18.9-liter depth-integrated sample was collected from each of the stations for chlorophyll *a* analysis. The maximum depths sampled were 29.3 meters at station 1, 34.7 meters at station 2, and 12.5 meters at station 3.

The sampling results are presented in full in Appendix D and are summarized in the following table.

A. SUMMARY OF PHYSICAL AND CHEMICAL CHARACTERISTICS FOR OXBOW RESERVOIR
STORET CODE 4106

| PARAMETER | 1ST SAMPLING ( 4/ 9/75) 3 SITES RANGE | MEAN | MEDIAN | 2ND SAMPLING ( 8/ 4/75) 3 SITES RANGE | MEAN | MEDIAN | 3RD SAMPLING ( 9/15/75) 3 SITES RANGE | MEAN | MEDIAN |
|---|---|---|---|---|---|---|---|---|---|
| TEMP (C) | 6.9 - 7.6 | 7.3 | 7.4 | 17.7 - 23.9 | 20.3 | 20.3 | 20.9 - 21.8 | 21.2 | 21.1 |
| DISS OXY (MG/L) | 15.2 - 15.6 | 15.5 | 15.6 | 2.8 - 9.2 | 6.0 | 5.8 | 4.0 - 8.6 | 4.9 | 4.6 |
| CNDCTVY (MCROMO) | 237. - 247. | 240. | 239. | 233. - 281. | 257. | 261. | 417. - 426. | 420. | 419. |
| PH (STAND UNITS) | 8.4 - 8.5 | 8.5 | 8.4 | 8.1 - 8.9 | 8.4 | 8.4 | 8.0 - 8.2 | 8.1 | 8.1 |
| TOT ALK (MG/L) | 129. - 134. | 131. | 131. | 95. - 102. | 99. | 101. | 151. - 171. | 162. | 166. |
| TOT P (MG/L) | 0.074 - 0.087 | 0.080 | 0.081 | 0.036 - 0.057 | 0.048 | 0.050 | 0.065 - 0.096 | 0.074 | 0.072 |
| ORTHO P (MG/L) | 0.024 - 0.051 | 0.036 | 0.037 | 0.012 - 0.043 | 0.032 | 0.034 | 0.063 - 0.067 | 0.065 | 0.064 |
| NO2+NO3 (MG/L) | 0.640 - 0.660 | 0.645 | 0.640 | 0.160 - 0.360 | 0.300 | 0.300 | 0.670 - 0.710 | 0.691 | 0.690 |
| AMMONIA (MG/L) | 0.040 - 0.060 | 0.052 | 0.050 | 0.020 - 0.040 | 0.027 | 0.030 | 0.020 - 0.070 | 0.023 | 0.020 |
| KJEL N (MG/L) | 0.300 - 0.500 | 0.385 | 0.400 | 0.200 - 0.300 | 0.250 | 0.250 | 0.200 - 0.500 | 0.269 | 0.250 |
| INORG N (MG/L) | 0.680 - 0.720 | 0.697 | 0.700 | 0.180 - 0.380 | 0.327 | 0.335 | 0.690 - 0.780 | 0.714 | 0.710 |
| TOTAL N (MG/L) | 0.940 - 1.150 | 1.030 | 1.040 | 0.460 - 0.660 | 0.550 | 0.540 | 0.890 - 1.210 | 0.960 | 0.940 |
| CHLRPYL A (UG/L) | 13.3 - 23.0 | 18.0 | 17.7 | 3.4 - 8.7 | 5.7 | 5.0 | 6.1 - 8.5 | 7.2 | 7.1 |
| SECCHI (METERS) | 0.5 - 0.8 | 0.7 | 0.7 | 0.4 - 3.2 | 2.3 | 3.2 | 2.4 - 3.4 | 2.7 | 2.4 |

B. Biological characteristics:

1. Phytoplankton -

| Sampling Date | Dominant Genera | Algal Units per ml |
|---|---|---|
| 04/09/75 | 1. Stephanodiscus sp. | 5,621 |
| | 2. Asterionella sp. | 300 |
| | 3. Melosira sp. | 257 |
| | 4. Surirella sp. | 43 |
| | 5. Chroomonas (?) sp. | 43 |
| | Other genera | 44 |
| | Total | 6,308 |
| 08/04/75 | 1. Fragilaria sp. | 433 |
| | 2. Stephanodiscus sp. | 31 |
| | Total | 464 |
| 09/15/75 | 1. Melosira sp. | 807 |
| | 2. Cyclotella sp. | 70 |
| | 3. Cryptomonas sp. | 35 |
| | Total | 912 |

2. Chlorophyll a -

| Sampling Date | Station Number | Chlorophyll a (µg/l) |
|---|---|---|
| 04/09/75 | 1 | 13.3 |
| | 2 | 23.0 |
| | 3 | 17.7 |
| 08/04/75 | 1 | 8.7 |
| | 2 | 5.0 |
| | 3 | 3.4 |
| 09/15/75 | 1 | 7.1 |
| | 2 | 8.5 |
| | 3 | 6.1 |

C. Limiting Nutrient Study:

The algal assay results are not considered representative of conditions in the reservoir at the time the sample was collected (09/15/75) due to significant changes in the nutrient levels in the sample during shipment from the field to the laboratory.

The reservoir data indicate phosphorus limitation in April and nitrogen limitation in August and September. The mean inorganic nitrogen/orthophosphorus ratios in April were 14/1 or greater at all stations and were 12/1 or less at all stations in August and September.

## IV. NUTRIENT LOADINGS
(See Appendix E for data)

For the determination of nutrient loadings, the Oregon National Guard collected monthly near-surface grab samples from each of the tributary sites indicated on the map (page v), except for the high runoff month of February when two samples were collected. Sampling was begun in October, 1974, and was completed in September, 1975.

Through an interagency agreement, stream flow estimates for the year of sampling and a "normalized" or average year were provided by the Oregon District Office of the U.S. Geological Survey for the tributary sites nearest the reservoir.

In this report, nutrient loads for sampled tributaries were determined by using a modification of a U.S. Geological Survey computer program for calculating stream loadings*. Nutrient loads for unsampled "minor tributaries and immediate drainage" ("ZZ" of U.S.G.S.) were estimated using the nutrient loads at station B-1, in $kg/km^2/year$, and multiplying by the ZZ area in $km^2$.

No known wastewater treatment plants directly impacted Oxbow Reservoir during the sampling year.

---

* See Working Paper No. 175.

A. Waste Sources:

1. Known municipal - None

2. Known industrial - None

B. Annual Total Phosphorus Loading - Average Year:

1. Inputs -

| Source | kg P/yr | % of total |
|---|---|---|
| a. Tributaries (non-point load) - | | |
| Snake River | 1,023,210 | 98.5 |
| Wildhorse River | 9,075 | 0.9 |
| b. Minor tributaries & immediate drainage (non-point load) - | 6,260 | 0.6 |
| c. Known municipal STP's - None | - | - |
| d. Septic tanks - None | - | - |
| e. Known industrial - None | - | - |
| f. Direct precipitation* - | 100 | < 0.1 |
| Total | 1,038,645 | 100.0 |

2. Outputs -

Reservoir outlet - Snake River 895,060

3. Net annual P accumulation - 143,585 kg.

* See Working Paper No. 175.

C. Annual Total Nitrogen Loading - Average Year:

1. Inputs -

| Source | kg N/yr | % of total |
|---|---|---|
| a. Tributaries (non-point load) - | | |
| Snake River | 18,218,785 | 98.6 |
| Wildhorse River | 151,760 | 0.8 |
| b. Minor tributaries & immediate drainage (non-point load) - | 103,570 | 0.6 |
| c. Known municipal STP's - None | - | - |
| d. Septic tanks - None | - | - |
| e. Known industrial - None | - | - |
| f. Direct precipitation* - | 6,110 | < 0.1 |
| Total | 18,480,225 | 100.0 |

2. Outputs -

Lake outlet - Snake River 25,222,530

3. Net annual N loss - 6,742,305 kg.

D. Non-point Nutrient Export by Subdrainage Area:

| Tributary | kg P/km²/yr | kg N/km²/yr |
|---|---|---|
| Snake River | 5 | 97 |
| Wildhorse River | 20 | 331 |

* See Working Paper No. 175.

E. Yearly Loads:

In the following table, the existing phosphorus loadings are compared to those proposed by Vollenweider (Vollenweider and Dillon, 1974). Essentially, his "dangerous" loading is one at which the receiving water would become eutrophic or remain eutrophic; his "permissible" loading is that which would result in the receiving water remaining oligotrophic or becoming oligotrophic if morphometry permitted. A mesotrophic loading would be considered one between "dangerous" and "permissible".

Note that Vollenweider's model may not be applicable to water bodies with short hydraulic retention times.

| | Total Phosphorus | | Total Nitrogen | |
|---|---|---|---|---|
| | Total | Accumulated | Total | Accumulated |
| grams/$m^2$/yr | 183.51 | 25.37 | 3,265.1 | loss* |

| Vollenweider phosphorus loadings ($g/m^2/yr$) based on mean depth and mean hydraulic retention time of Oxbow Reservoir: | |
|---|---|
| "Dangerous" (eutrophic loading) | 9.00 |
| "Permissible" (oligotrophic loading) | 4.50 |

* There was an apparent loss of nitrogen during the sampling year. This may have been due to nitrogen fixation in the reservoir, solubilization of previously sedimented nitrogen, recharge with nitrogen-rich ground water, unsampled point sources discharging directly to the reservoir, or (probably) insufficient outlet sampling in relation to the very short hydraulic retention time of the reservoir. Whatever the cause, a similar nitrogen loss has occurred at Shagawa Lake, Minnesota, which has been intensively studied by EPA's former National Eutrophication and Lake Restoration Branch (Malueg et al., 1975).

## V. LITERATURE REVIEWED

Malueg, Kenneth W., D. Phillips Larsen, Donald W. Schults, and Howard T. Mercier; 1975. A six-year water, phosphorus, and nitrogen budget for Shagawa Lake, Minnesota. Jour. Environ. Qual., vol. 4, no. 2, pp. 236-242.

Martin, R. A. R., and Ronald L. Hanson, 1966. Reservoirs in the United States. Water Supply Paper No. 1838, U.S. Geol. Surv., Wash., DC.

McHugh, Robert A., 1972. Interim study of some physical, chemical, and biological properties of selected Oregon lakes. OR Dept. of Environ. Qual., Portland.

Vollenweider, R. A., and P. J. Dillon, 1974. The application of the phosphorus loading concept to eutrophication research. Natl. Res. Council of Canada Publ. No. 13690, Canada Centre for Inland Waters, Burlington, Ontario.

# VI. APPENDICES

## APPENDIX A

## LAKE RANKINGS

LAKE DATA TO BE USED IN RANKINGS

Reproduced from best available copy.

| LAKE CODE | LAKE NAME | MEDIAN TOTAL P | MEDIAN INORG N | 500-MEAN SEC | MEAN CHLORA | 15-MIN DO | MEDIAN DISS ORTHO P |
|---|---|---|---|---|---|---|---|
| 1601 | AMERICAN FALLS RESERVOIR | 0.105 | 0.080 | 463.800 | 15.379 | 14.700 | 0.035 |
| 1602 | CASCADE LAKE | 0.032 | 0.060 | 415.067 | 8.081 | 14.600 | 0.009 |
| 1603 | LAKE COEUR D'ALENE | 0.017 | 0.040 | 380.348 | 10.391 | 12.200 | 0.005 |
| 1604 | DWORSHAK RESERVOIR | 0.010 | 0.080 | 401.866 | 2.420 | 7.400 | 0.009 |
| 1605 | HAUSER | 0.028 | 0.075 | 366.286 | 11.112 | 14.800 | 0.013 |
| 1606 | HAYDEN LAKE | 0.010 | 0.040 | 243.500 | 2.787 | 11.800 | 0.003 |
| 1607 | ISLAND PARK RESERVOIR | 0.034 | 0.050 | 391.778 | 9.322 | 12.800 | 0.012 |
| 1608 | LAKE LOWELL | 0.070 | 0.070 | 477.111 | 25.389 | 14.600 | 0.015 |
| 1609 | MAGIC RESERVOIR | 0.062 | 0.130 | 400.750 | 7.322 | 14.700 | 0.020 |
| 1610 | PALISADES RESERVOIR | 0.024 | 0.080 | 345.428 | 2.067 | 12.800 | 0.007 |
| 1611 | LOWER PAYETTE | 0.013 | 0.060 | 234.000 | 4.600 | 9.600 | 0.007 |
| 1612 | LOWER TWIN LAKES | 0.016 | 0.050 | 370.000 | 2.318 | 13.600 | 0.009 |
| 1613 | UPPER TWIN LAKES | 0.017 | 0.045 | 369.143 | 34.962 | 8.200 | 0.004 |
| 4101 | BROWNLEE RESERVOIR | 0.079 | 0.560 | 428.133 | 16.207 | 14.500 | 0.043 |
| 4102 | DIAMOND LAKE | 0.028 | 0.040 | 294.500 | 7.300 | 6.800 | 0.011 |
| 4103 | HELLS CANYON RESERVOIR | 0.068 | 0.640 | 429.111 | 18.722 | 12.400 | 0.045 |
| 4104 | HILLS CREEK RESERVOIR | 0.030 | 0.060 | 435.200 | 2.333 | 7.400 | 0.027 |
| 4105 | OWYHEE | 0.095 | 0.425 | 480.417 | 3.350 | 13.200 | 0.064 |
| 4106 | OXBOW RESERVOIR | 0.071 | 0.690 | 425.555 | 10.311 | 12.200 | 0.040 |
| 4107 | SUTTLE LAKE | 0.031 | 0.040 | 95.000 | 9.167 | 6.800 | 0.020 |
| 4108 | WALDO LAKE | 0.005 | 0.040 | -100.000 | 0.350 | 6.800 | 0.006 |

PERCENT OF LAKES WITH HIGHER VALUES (NUMBER OF LAKES WITH HIGHER VALUES)

| LAKE CODE | LAKE NAME | MEDIAN TOTAL P | MEDIAN INORG N | 500-MEAN SEC | MEAN CHLORA | 15-MIN DO | MEDIAN DISS ORTHO P | INDEX NO |
|---|---|---|---|---|---|---|---|---|
| 1601 | AMERICAN FALLS RESERVOIR | 0 ( 0) | 30 ( 5) | 10 ( 2) | 20 ( 4) | 13 ( 2) | 20 ( 4) | 93 |
| 1602 | CASCADE LAKE | 45 ( 9) | 58 ( 11) | 35 ( 7) | 50 ( 10) | 3 ( 0) | 65 ( 12) | 256 |
| 1603 | LAKE COEUR D'ALENE | 75 ( 15) | 93 ( 17) | 55 ( 11) | 30 ( 6) | 58 ( 11) | 90 ( 18) | 401 |
| 1604 | DWORSHAK RESERVOIR | 93 ( 18) | 30 ( 5) | 40 ( 8) | 80 ( 16) | 83 ( 16) | 65 ( 12) | 391 |
| 1605 | HAUSER | 58 ( 11) | 40 ( 8) | 70 ( 14) | 25 ( 5) | 3 ( 0) | 45 ( 9) | 241 |
| 1606 | HAYDEN LAKE | 93 ( 18) | 93 ( 17) | 85 ( 17) | 75 ( 15) | 65 ( 13) | 100 ( 20) | 511 |
| 1607 | ISLAND PARK RESERVOIR | 40 ( 8) | 68 ( 13) | 50 ( 10) | 40 ( 8) | 43 ( 8) | 50 ( 10) | 291 |
| 1608 | LAKE LOWELL | 20 ( 4) | 45 ( 9) | 5 ( 1) | 5 ( 1) | 20 ( 4) | 40 ( 8) | 135 |
| 1609 | MAGIC RESERVOIR | 30 ( 6) | 20 ( 4) | 45 ( 9) | 55 ( 11) | 13 ( 2) | 30 ( 6) | 193 |
| 1610 | PALISADES RESERVOIR | 65 ( 13) | 30 ( 5) | 75 ( 15) | 95 ( 19) | 43 ( 8) | 80 ( 16) | 388 |
| 1611 | LOWER PAYETTE | 85 ( 17) | 58 ( 11) | 90 ( 18) | 65 ( 13) | 70 ( 14) | 75 ( 15) | 443 |
| 1612 | LOWER TWIN LAKES | 80 ( 16) | 68 ( 13) | 60 ( 12) | 90 ( 18) | 30 ( 6) | 65 ( 12) | 393 |
| 1613 | UPPER TWIN LAKES | 70 ( 14) | 75 ( 15) | 65 ( 13) | 0 ( 0) | 75 ( 15) | 95 ( 19) | 380 |
| 4101 | BROWNLEE RESERVOIR | 10 ( 2) | 10 ( 2) | 25 ( 5) | 15 ( 3) | 25 ( 5) | 10 ( 2) | 95 |
| 4102 | DIAMOND LAKE | 58 ( 11) | 93 ( 17) | 80 ( 16) | 60 ( 12) | 95 ( 18) | 55 ( 11) | 441 |
| 4103 | HELLS CANYON RESERVOIR | 25 ( 5) | 5 ( 1) | 20 ( 4) | 10 ( 2) | 50 ( 10) | 5 ( 1) | 115 |
| 4104 | HILLS CREEK RESERVOIR | 35 ( 7) | 50 ( 10) | 15 ( 3) | 85 ( 17) | 83 ( 16) | 25 ( 5) | 293 |
| 4105 | OWYHEE | 5 ( 1) | 15 ( 3) | 0 ( 0) | 70 ( 14) | 35 ( 7) | 0 ( 0) | 125 |
| 4106 | OXBOW RESERVOIR | 15 ( 3) | 0 ( 0) | 30 ( 6) | 35 ( 7) | 58 ( 11) | 15 ( 3) | 153 |
| 4107 | SUTTLE LAKE | 50 ( 10) | 80 ( 16) | 95 ( 19) | 45 ( 9) | 95 ( 18) | 35 ( 7) | 400 |
| 4108 | WALDO LAKE | 100 ( 20) | 93 ( 17) | 100 ( 20) | 100 ( 20) | 95 ( 18) | 85 ( 17) | 573 |

LAKES RANKED BY INDEX NOS.

| RANK | LAKE CODE | LAKE NAME | INDEX NO |
|---|---|---|---|
| 1 | 4108 | WALDO LAKE | 573 |
| 2 | 1606 | HAYDEN LAKE | 511 |
| 3 | 1611 | LOWER PAYETTE | 443 |
| 4 | 4102 | DIAMOND LAKE | 441 |
| 5 | 1603 | LAKE COEUR D'ALENE | 401 |
| 6 | 4107 | SUTTLE LAKE | 400 |
| 7 | 1612 | LOWER TWIN LAKES | 393 |
| 8 | 1604 | DWORSHAK RESERVOIR | 391 |
| 9 | 1610 | PALISADES RESERVOIR | 388 |
| 10 | 1613 | UPPER TWIN LAKES | 380 |
| 11 | 4104 | HILLS CREEK RESERVOIR | 293 |
| 12 | 1607 | ISLAND PARK RESERVOIR | 291 |
| 13 | 1602 | CASCADE LAKE | 256 |
| 14 | 1605 | HAUSER | 241 |
| 15 | 1609 | MAGIC RESERVOIR | 193 |
| 16 | 4106 | OXBOW RESERVOIR | 153 |
| 17 | 1608 | LAKE LOWELL | 135 |
| 18 | 4105 | OWYHEE | 125 |
| 19 | 4103 | HELLS CANYON RESERVOIR | 115 |
| 20 | 4101 | BROWNLEE RESERVOIR | 95 |
| 21 | 1601 | AMERICAN FALLS RESERVOIR | 93 |

# APPENDIX B

## CONVERSION FACTORS

## CONVERSION FACTORS

Hectares x 2.471 = acres

Kilometers x 0.6214 = miles

Meters x 3.281 = feet

Cubic meters x $8.107 \times 10^{-4}$ = acre/feet

Square kilometers x 0.3861 = square miles

Cubic meters/sec x 35.315 - cubic feet/sec

Centimeters x 0.3937 = inches

Kilograms x 2.205 = pounds

Kilograms/square kilometer x 5.711 = lbs/square mile

# APPENDIX C

## TRIBUTARY FLOW DATA

TRIBUTARY FLOW INFORMATION FOR OREGON 08/11/76

LAKE CODE 4106 OX BOW RESERVOIR

TOTAL DRAINAGE AREA OF LAKE(SQ KM) 188810.9

| TRIBUTARY | SUB-DRAINAGE AREA(SQ KM) | NORMALIZED FLOWS(CMS) JAN | FEB | MAR | APR | MAY | JUN | JUL | AUG | SEP | OCT | NOV | DEC | MEAN |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 4106A1 | 188810.9 | 597.49 | 713.58 | 676.77 | 716.42 | 634.30 | 620.14 | 336.97 | 325.64 | 396.44 | 447.41 | 444.57 | 501.21 | 532.66 |
| 4106A2 | 188033.9 | 594.65 | 710.75 | 671.11 | 707.92 | 614.48 | 600.32 | 331.31 | 322.81 | 396.44 | 444.57 | 441.74 | 498.38 | 526.27 |
| 4106B1 | 458.4 | 2.27 | 2.12 | 5.10 | 6.51 | 15.57 | 15.57 | 4.25 | 1.42 | 0.99 | 1.42 | 2.27 | 1.42 | 4.92 |
| 4106ZZ | 89.1 | 0.42 | 0.51 | 0.68 | 0.79 | 2.41 | 1.42 | 0.68 | 0.23 | 0.17 | 0.20 | 0.23 | 0.23 | 0.66 |

SUMMARY

TOTAL DRAINAGE AREA OF LAKE = 188810.9
SUM OF SUB-DRAINAGE AREAS = 188581.4

TOTAL FLOW IN = 6401.27
TOTAL FLOW OUT = 6410.92

MEAN MONTHLY FLOWS AND DAILY FLOWS(CMS)

| TRIBUTARY | MONTH | YEAR | MEAN FLOW | DAY | FLOW | DAY | FLOW | DAY | FLOW |
|---|---|---|---|---|---|---|---|---|---|
| 4106A1 | 10 | 74 | 501.208 | 19 | 410.594 | | | | |
| | 11 | 74 | 495.545 | 23 | 521.030 | | | | |
| | 12 | 74 | 523.862 | 14 | 538.020 | | | | |
| | 1 | 75 | 614.476 | 4 | 620.139 | 25 | 656.951 | | |
| | 2 | 75 | 719.248 | 9 | 569.168 | 22 | 894.812 | | |
| | 3 | 75 | 877.822 | 9 | 897.644 | | | | |
| | 4 | 75 | 1268.595 | 13 | 1155.327 | | | | |
| | 5 | 75 | 1347.882 | 17 | 82.119 | | | | |
| | 6 | 75 | 733.406 | 7 | 744.733 | | | | |
| | 7 | 75 | 413.426 | | | | | | |
| | 8 | 75 | 328.475 | 9 | 311.485 | | | | |
| | 9 | 75 | 444.574 | 28 | 430.416 | | | | |
| 4106A2 | 10 | 74 | 498.376 | 19 | 410.594 | | | | |
| | 11 | 74 | 492.713 | 23 | 521.030 | | | | |
| | 12 | 74 | 523.862 | 14 | 538.020 | | | | |
| | 1 | 75 | 614.476 | 4 | 617.307 | 25 | 656.951 | | |
| | 2 | 75 | 719.248 | 9 | 569.168 | 22 | 891.980 | | |
| | 3 | 75 | 872.159 | 9 | 889.149 | | | | |
| | 4 | 75 | 1260.100 | 13 | 1149.664 | | | | |
| | 5 | 75 | 1328.060 | 17 | 51.537 | | | | |
| | 6 | 75 | 719.248 | 7 | 722.080 | | | | |
| | 7 | 75 | 407.762 | | | | | | |
| | 8 | 75 | 325.644 | 9 | 308.654 | | | | |
| | 9 | 75 | 444.574 | 28 | 430.416 | | | | |

12

TRIBUTARY FLOW INFORMATION FOR OREGON 08/11/76

LAKE CODE 4106 OX BOW RESERVOIR

MEAN MONTHLY FLOWS AND DAILY FLOWS(CMS)

| TRIBUTARY | MONTH | YEAR | MEAN FLOW | DAY | FLOW | DAY | FLOW | DAY | FLOW |
|---|---|---|---|---|---|---|---|---|---|
| 4106d1 | 10 | 74 | 0.850 | 19 | 0.906 | | | | |
| | 11 | 74 | 0.566 | 23 | 0.623 | | | | |
| | 12 | 74 | 0.623 | 14 | 0.623 | | | | |
| | 1 | 75 | 0.708 | 4 | 0.623 | 25 | 0.850 | | |
| | 2 | 75 | 1.416 | 9 | 0.991 | 22 | 1.189 | | |
| | 3 | 75 | 5.380 | 9 | 8.495 | | | | |
| | 4 | 75 | 5.947 | 13 | 5.663 | | | | |
| | 5 | 75 | 17.556 | 17 | 25.485 | | | | |
| | 6 | 75 | 12.743 | 7 | 18.406 | | | | |
| | 7 | 75 | 4.531 | | | | | | |
| | 8 | 75 | 1.076 | 9 | 0.708 | | | | |
| | 9 | 75 | 0.793 | 28 | 0.566 | | | | |

# APPENDIX D

## PHYSICAL and CHEMICAL DATA

STORET RETRIEVAL DATE 76/08/12

410601
44 58 07.0 116 50 02.0 3
OXBOW RESERVOIR
41001 OREGON

130892

11EPALES 2111202
0064 FEET DEPTH CLASS 00

| DATE FROM TO | TIME OF DAY | DEPTH FEET | 00010 WATER TEMP CENT | 00300 DO MG/L | 00077 TRANSP SECCHI INCHES | 00094 CNDUCTVY FIELD MICROMHO | 00400 PH SU | 00410 T ALK CACO3 MG/L | 00610 NH3-N TOTAL MG/L | 00625 TOT KJEL N MG/L | 00630 NO2&NO3 N-TOTAL MG/L | 00671 PHOS-DIS ORTHO MG/L P |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 75/04/09 | 10 00 | 0000 | 6.9 | 15.4 | 27 | 247 | 8.50 | 132 | 0.050 | 0.400 | 0.650 | 0.033 |
| | 10 00 | 0005 | 7.1 | 15.6 | | 239 | 8.45 | 131 | 0.040 | 0.400 | 0.640 | 0.037 |
| | 10 00 | 0015 | 7.1 | 15.2 | | 239 | 8.40 | 130 | 0.050 | 0.400 | 0.640 | 0.042 |
| | 10 00 | 0035 | 7.0 | 15.6 | | 240 | 8.45 | 129 | 0.040 | 0.400 | 0.640 | 0.039 |
| | 10 00 | 0060 | 7.0 | 15.4 | | 243 | 8.45 | 129 | 0.050 | 0.400 | 0.640 | 0.040 |
| 75/08/04 | 09 00 | 0000 | 23.9 | 9.2 | 126 | 281 | 8.90 | 101 | 0.020 | 0.300 | 0.160 | 0.012 |
| | 09 00 | 0005 | 22.2 | 7.6 | | 275 | 8.70 | 102 | 0.030 | 0.300 | 0.240 | 0.019 |
| | 09 00 | 0015 | 21.1 | 5.6 | | 269 | 8.40 | 102 | 0.030 | 0.300 | 0.280 | 0.027 |
| | 09 00 | 0035 | 20.6 | 6.0 | | 265 | 8.35 | 101 | 0.020 | 0.200 | 0.300 | 0.029 |
| | 09 00 | 0070 | 20.0 | 4.8 | | 249 | 8.15 | 97 | 0.030 | 0.200 | 0.320 | 0.034 |
| | 09 00 | 0096 | 19.2 | 5.8 | | 241 | 8.65 | 96 | 0.030 | 0.300 | 0.350 | 0.037 |
| 75/09/15 | 15 00 | 0000 | 21.6 | 5.6 | 96 | 426 | 8.20 | 167 | 0.020K | 0.200 | 0.690 | 0.067 |
| | 15 00 | 0005 | 21.0 | 8.6 | | 419 | 8.10 | 166 | 0.020K | 0.200 | 0.700 | 0.065 |
| | 15 00 | 0015 | 21.0 | 4.5 | | 417 | 8.10 | 165 | 0.020K | 0.200 | 0.700 | 0.066 |
| | 15 00 | 0035 | 20.9 | 4.2 | | 417 | 8.10 | 165 | 0.020K | 0.200 | 0.710 | 0.066 |
| | 15 00 | 0065 | 20.9 | 4.3 | | 417 | 8.10 | 166 | 0.070 | 0.500 | 0.710 | 0.066 |
| | 15 00 | 0096 | 20.9 | 4.0 | | 420 | 8.10 | 166 | 0.020K | 0.200 | 0.710 | 0.067 |

| DATE FROM TO | TIME OF DAY | DEPTH FEET | 00665 PHOS-TOT MG/L P | 32217 CHLRPHYL A UG/L | 00031 INCDT LT REMNING PERCENT |
|---|---|---|---|---|---|
| 75/04/09 | 10 00 | 0000 | 0.087 | 13.3 | |
| | 10 00 | 0005 | 0.084 | | |
| | 10 00 | 0015 | 0.074 | | |
| | 10 00 | 0035 | 0.079 | | |
| | 10 00 | 0060 | 0.079 | | |
| 75/08/04 | 09 00 | 0000 | 0.036 | 8.7 | |
| | 09 00 | 0005 | 0.040 | | |
| | 09 00 | 0015 | 0.040 | | |
| | 09 00 | 0035 | 0.042 | | |
| | 09 00 | 0070 | 0.051 | | |
| | 09 00 | 0096 | 0.055 | | |
| 75/09/15 | 15 00 | 0000 | 0.073 | 7.1 | |
| | 15 00 | 0005 | 0.069 | | |
| | 15 00 | 0015 | 0.071 | | |
| | 15 00 | 0035 | 0.072 | | |
| | 15 00 | 0065 | 0.096 | | |
| | 15 00 | 0096 | 0.085 | | |

K VALUE KNOWN TO BE
LESS THAN INDICATED

STORET RETRIEVAL DATE 76/08/12

410602
44 56 10.0 116 49 58.0 3
OXBOW RESERVOIR
41001 OREGON

130892

11EPALES 2111202
0094 FEET DEPTH CLASS 00

| DATE FROM TO | TIME OF DAY | DEPTH FEET | 00010 WATER TEMP CENT | 00300 DO MG/L | 00077 TRANSP SECCHI INCHES | 00094 CNDUCTVY FIELD MICROMHO | 00400 PH SU | 00410 T ALK CACO3 MG/L | 00610 NH3-N TOTAL MG/L | 00625 TOT KJEL N MG/L | 00630 NO2&NO3 N-TOTAL MG/L | 00671 PHOS-DIS ORTHO MG/L P |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 75/04/09 | 10 | 25 0000 | 7.2 | 15.6 | 30 | 239 | 8.50 | 129 | 0.060 | 0.300 | 0.650 | 0.041 |
| | 10 | 25 0005 | 7.5 | 15.6 | | 239 | 8.50 | 130 | 0.050 | 0.400 | 0.640 | 0.044 |
| | 10 | 25 0015 | 7.5 | 15.6 | | 238 | 8.50 | 129 | 0.060 | 0.400 | 0.650 | 0.051 |
| | 10 | 25 0050 | 7.4 | 15.4 | | 239 | 8.45 | 134 | 0.060 | 0.300 | 0.660 | 0.033 |
| | 10 | 25 0090 | 7.4 | 15.6 | | 239 | 8.45 | 132 | 0.050 | 0.400 | 0.650 | 0.028 |
| 75/08/04 | 09 | 30 0000 | 21.7 | | 126 | 273 | 8.60 | 101 | 0.030 | 0.300 | 0.280 | 0.034 |
| | 09 | 30 0005 | 21.7 | 8.0 | | 275 | 8.60 | 101 | 0.040 | 0.300 | 0.270 | 0.037 |
| | 09 | 30 0015 | 21.0 | 4.4 | | 270 | 8.40 | 100 | 0.040 | 0.200K | 0.300 | 0.039 |
| | 09 | 30 0030 | 20.8 | | | 265 | 8.35 | 97 | 0.030 | 0.200 | 0.330 | 0.036 |
| | 09 | 30 0060 | 19.9 | 3.0 | | 257 | 8.20 | 99 | 0.030 | 0.200 | 0.330 | 0.035 |
| | 09 | 30 0090 | 19.5 | 2.8 | | 247 | 8.10 | 99 | 0.020 | 0.200 | 0.340 | 0.035 |
| | 09 | 30 0114 | 19.5 | 7.0 | | 247 | 8.55 | 101 | 0.030 | 0.200 | 0.300 | 0.030 |
| 75/09/15 | 15 | 20 0000 | 21.8 | 5.3 | 96 | 423 | 8.20 | 165 | 0.020K | 0.200 | 0.690 | 0.063 |
| | 15 | 20 0005 | 21.5 | 5.2 | | 423 | 8.20 | 166 | 0.020K | 0.200 | 0.690 | 0.063 |
| | 15 | 20 0015 | 21.3 | 5.1 | | 419 | 8.20 | 169 | 0.020K | 0.200 | 0.690 | 0.063 |
| | 15 | 20 0035 | 21.1 | 4.9 | | 418 | 8.10 | 169 | 0.020K | 0.300 | 0.670 | 0.064 |
| | 15 | 20 0070 | 21.1 | 4.6 | | 418 | 8.10 | 171 | 0.020K | 0.300 | 0.690 | 0.065 |
| | 15 | 20 0101 | 21.1 | 4.5 | | 419 | 8.00 | 154 | 0.020K | 0.400 | 0.690 | 0.067 |

| DATE FROM TO | TIME OF DAY | DEPTH FEET | 00665 PHOS-TOT MG/L P | 32217 CHLRPHYL A UG/L | 00031 INCDT LT REMNING PERCENT |
|---|---|---|---|---|---|
| 75/04/09 | 10 | 25 0000 | 0.081 | 23.0 | |
| | 10 | 25 0005 | 0.082 | | |
| | 10 | 25 0015 | 0.085 | | |
| | 10 | 25 0050 | 0.081 | | |
| | 10 | 25 0090 | 0.081 | | |
| 75/08/04 | 09 | 30 0000 | 0.042 | 5.0 | |
| | 09 | 30 0005 | 0.043 | | |
| | 09 | 30 0015 | 0.041 | | |
| | 09 | 30 0030 | 0.055 | | |
| | 09 | 30 0060 | 0.051 | | |
| | 09 | 30 0090 | 0.051 | | |
| | 09 | 30 0114 | 0.057 | | |
| 75/09/15 | 15 | 20 0000 | 0.071 | 8.5 | |
| | 15 | 20 0005 | 0.073 | | |
| | 15 | 20 0015 | 0.076 | | |
| | 15 | 20 0035 | 0.074 | | |
| | 15 | 20 0070 | 0.078 | | |
| | 15 | 20 0101 | 0.085 | | |

K VALUE KNOWN TO BE
LESS THAN INDICATED

STORET RETRIEVAL DATE 76/08/12

410603
44 51 23.0 116 53 25.0 3
OXBOW RESERVOIR
41001 OREGON

130892

11EPALES 2111202
0019 FEET DEPTH CLASS 00

| DATE FROM TO | TIME OF DAY | DEPTH FEET | 00010 WATER TEMP CENT | 00300 DO MG/L | 00077 TRANSP SECCHI INCHES | 00094 CNDUCTVY FIELD MICROMHO | 00400 PH SU | 00410 T ALK CACO3 MG/L | 00610 NH3-N TOTAL MG/L | 00625 TOT KJEL N MG/L | 00630 NO2&NO3 N-TOTAL MG/L | 00671 PHOS-DIS ORTHO MG/L P |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 75/04/09 | 11 00 | 0000 | 7.5 | 15.6 | 20 | 237 | 8.45 | 131 | 0.050 | 0.500 | 0.650 | 0.024 |
| | 11 00 | 0005 | 7.6 | 15.2 | | 237 | 8.45 | 132 | 0.060 | 0.300 | 0.640 | 0.029 |
| | 11 00 | 0015 | 7.4 | 15.2 | | 238 | 8.50 | 132 | 0.050 | 0.400 | 0.640 | 0.026 |
| 75/08/04 | 10 05 | 0000 | 17.7 | 8.8 | 17 | 233 | 8.55 | 102 | 0.020 | 0.300 | 0.280 | 0.025 |
| | 10 05 | 0005 | 17.8 | 5.4 | | 235 | 8.20 | 97 | 0.020 | 0.300 | 0.360 | 0.042 |
| | 10 05 | 0013 | 17.8 | 5.8 | | 233 | 8.10 | 95 | 0.020 | 0.200 | 0.360 | 0.043 |
| 75/09/15 | 16 40 | 0000 | 21.2 | 4.6 | 132 | 420 | 8.15 | 153 | 0.020K | 0.300 | 0.680 | 0.063 |
| | 16 40 | 0005 | 21.2 | 4.6 | | 420 | 8.20 | 154 | 0.020K | 0.300 | 0.680 | 0.063 |
| | 16 40 | 0017 | 21.2 | 4.6 | | 421 | 8.20 | 151 | 0.020K | 0.300 | 0.680 | 0.064 |
| | 16 40 | 0041 | 21.2 | 4.5 | | 419 | 8.10 | 151 | 0.020K | 0.300 | 0.680 | 0.063 |

| DATE FROM TO | TIME OF DAY | DEPTH FEET | 00665 PHOS-TOT MG/L P | 32217 CHLRPHYL A UG/L | 00031 INCDT LT REMNING PERCENT |
|---|---|---|---|---|---|
| 75/04/09 | 11 00 | 0000 | 0.076 | 17.7 | |
| | 11 00 | 0005 | 0.075 | | |
| | 11 00 | 0015 | 0.079 | | |
| 75/08/04 | 10 05 | 0000 | 0.049 | 3.4 | |
| | 10 05 | 0005 | 0.053 | | |
| | 10 05 | 0013 | 0.056 | | |
| 75/09/15 | 16 40 | 0000 | 0.068 | 6.1 | |
| | 16 40 | 0005 | 0.065 | | |
| | 16 40 | 0017 | 0.065 | | |
| | 16 40 | 0041 | 0.067 | | |

K VALUE KNOWN TO BE
LESS THAN INDICATED

# APPENDIX E

# TRIBUTARY DATA

STORET RETRIEVAL DATE 76/08/12

4106A1
44 58 30.0 116 51 15.0 4
SNAKE RIVER
41 15 COPPERFIELD
O/OX BOW RESERVOIR 130892
BRDG BELOW OXBOW POWERHOUSE
11EPALES 2111204
0000 FEET DEPTH CLASS 00

| DATE FROM TO | TIME OF DAY | DEPTH FEET | 00630 NO2&NO3 N-TOTAL MG/L | 00625 TOT KJEL N MG/L | 00610 NH3-N TOTAL MG/L | 00671 PHOS-DIS ORTHO MG/L P | 00665 PHOS-TOT MG/L P |
|---|---|---|---|---|---|---|---|
| 74/10/19 | 12 50 | | 0.800 | 1.400 | 0.055 | 0.065 | 0.075 |
| 74/11/23 | 12 30 | | 0.960 | 0.400 | 0.035 | 0.050 | 0.050 |
| 75/01/25 | 14 22 | | 0.810 | 0.500 | 0.056 | 0.016 | 0.030 |
| 75/02/09 | 14 50 | | 0.750 | 1.100 | 0.072 | 0.024 | 0.050 |
| 75/02/22 | 15 35 | | 0.720 | 0.400 | 0.044 | 0.016 | 0.040 |
| 75/05/17 | 14 00 | | 0.070 | 0.750 | 0.030 | 0.010 | 0.090 |
| 75/06/07 | 13 00 | | 0.055 | 1.300 | 0.030 | 0.010 | 0.060 |
| 75/08/09 | | | 0.300 | 1.850 | 0.050 | 0.030 | 0.060 |
| 75/09/28 | 11 00 | | 0.620 | 0.500 | 0.025 | 0.050 | 0.070 |

STORET RETRIEVAL DATE 76/08/12

4106A2
44 50 45.0 116 53 45.0 4
SNAKE RIVER
41 15 COPPERFIELD
T/OX BOW RESERVOIR 130892
PVD HWY BRDG .4 MI N OF BROWNLEE DAM
11EPALES 2111204
0000 FEET DEPTH CLASS 00

| DATE FROM TO | TIME OF DAY | DEPTH FEET | 00630 NO2&NO3 N-TOTAL MG/L | 00625 TOT KJEL N MG/L | 00610 NH3-N TOTAL MG/L | 00671 PHOS-DIS ORTHO MG/L P | 00665 PHOS-TOT MG/L P |
|---|---|---|---|---|---|---|---|
| 74/10/19 | 14 30 | | 0.800 | 0.600 | 0.065 | 0.060 | 0.070 |
| 74/11/23 | 12 00 | | 0.950 | 0.400 | 0.075 | 0.045 | 0.045 |
| 74/12/14 | 13 30 | | 0.990 | 0.700 | 0.060 | 0.045 | 0.050 |
| 75/02/09 | 11 15 | | 0.760 | 0.600 | 0.056 | 0.016 | 0.050 |
| 75/02/22 | 12 00 | | 0.720 | 0.400 | 0.056 | 0.016 | 0.040 |
| 75/03/09 | 12 00 | | 0.825 | 1.100 | 0.044 | 0.024 | 0.070 |
| 75/04/13 | 10 55 | | 0.500 | 1.850 | 0.055 | 0.020 | 0.080 |
| 75/05/17 | 11 35 | | 0.060 | 1.630 | 0.045 | 0.015 | 0.080 |
| 75/06/07 | 12 25 | | 0.050 | 0.550 | 0.040 | 0.010 | 0.050 |
| 75/09/28 | | | 0.640 | 1.000 | 0.045 | 0.060 | 0.090 |

STORET RETRIEVAL DATE 76/08/12

410681
44 51 15.0 116 53 30.0 4
WILDHORSE RIVER
41 15 COPPERFIELD
T/OX BOW RESERVOIR 130892
BRDG .4 MI N OF ADAMS/WASH CO LINE
11EPALES 2111204
0000 FEET DEPTH CLASS 00

| DATE FROM TO | TIME OF DAY | DEPTH FEET | 00630 NO2&NO3 N-TOTAL MG/L | 00625 TOT KJEL N MG/L | 00610 NH3-N TOTAL MG/L | 00671 PHOS-DIS ORTHO MG/L P | 00665 PHOS-TOT MG/L P |
|---|---|---|---|---|---|---|---|
| 74/10/14 | 14 15 | | 0.040 | 0.600 | 0.025 | 0.015 | 0.020 |
| 74/11/23 | 11 45 | | 0.016 | 0.200 | 0.030 | 0.015 | 0.020 |
| 74/12/14 | 13 06 | | 0.056 | 0.300 | 0.010 | 0.015 | 0.020 |
| 75/01/04 | 11 00 | | 0.096 | 0.700 | 0.020 | 0.028 | 0.040 |
| 75/01/25 | 10 55 | | 0.088 | 0.700 | 0.016 | 0.016 | 0.020 |
| 75/02/09 | 11 00 | | 0.152 | 0.600 | 0.040 | 0.016 | 0.030 |
| 75/02/22 | 12 05 | | 0.232 | 0.100 | 0.012 | 0.016 | 0.020 |
| 75/03/09 | 12 04 | | 0.297 | 0.700 | 0.032 | 0.024 | 0.070 |
| 75/04/13 | 10 45 | | 0.105 | 1.050 | 0.020 | 0.025 | 0.080 |
| 75/05/17 | 11 30 | | 0.070 | 0.850 | 0.080 | 0.025 | |
| 75/06/07 | 12 20 | | 0.080 | 1.050 | 0.015 | 0.015 | 0.050 |
| 75/09/28 | | | 0.045 | 0.600 | 0.015 | 0.010 | 0.050 |